

## The master method

The master method applies to recurrences of the form

$$T(n) = a T(n/b) + f(n),$$

where  $a \geq 1$ ,  $b > 1$ , and  $f$  is asymptotically positive.

Slides courtesy of Charles Leiserson with small changes by Carola Wenk  
Fall 2009 CS 2233 Discr. Math. Structures

1

## Master theorem

$$T(n) = a T(n/b) + f(n)$$

**CASE 1:**  $f(n) \in O(n^{\log_b a - \varepsilon})$   
 $\Rightarrow T(n) \in \Theta(n^{\log_b a})$ .

**CASE 2:**  $f(n) \in \Theta(n^{\log_b a})$   
 $\Rightarrow T(n) \in \Theta(n^{\log_b a} \log n)$ .

**CASE 3:**  $f(n) \in \Omega(n^{\log_b a + \varepsilon})$  and  $af(n/b) \leq c f(n)$   
for some  $c < 1$   
 $\Rightarrow T(n) \in \Theta(f(n))$ .

**Merge sort:**  $a = 2$ ,  $b = 2 \Rightarrow n^{\log_b a} = n$   
 $\Rightarrow$  CASE 2  $\Rightarrow T(n) \in \Theta(n \log n)$ .

Fall 2009 CS 2233 Discr. Math. Structures

2

## Three common cases

Compare  $f(n)$  with  $n^{\log_b a}$ :

1.  $f(n) \in O(n^{\log_b a - \varepsilon})$  for some constant  $\varepsilon > 0$ .
  - $f(n)$  grows polynomially slower than  $n^{\log_b a}$  (by an  $n^\varepsilon$  factor).

**Solution:**  $T(n) \in \Theta(n^{\log_b a})$ .

2.  $f(n) \in \Theta(n^{\log_b a})$ .
  - $f(n)$  and  $n^{\log_b a}$  grow at similar rates.

**Solution:**  $T(n) \in \Theta(n^{\log_b a} \log n)$ .

Fall 2009

CS 2233 Discr. Math. Structures

3

## Three common cases (cont.)

Compare  $f(n)$  with  $n^{\log_b a}$ :

3.  $f(n) \in \Omega(n^{\log_b a + \varepsilon})$  for some constant  $\varepsilon > 0$ .
    - $f(n)$  grows polynomially faster than  $n^{\log_b a}$  (by an  $n^\varepsilon$  factor),  
and  $f(n)$  satisfies the **regularity condition** that  $af(n/b) \leq c f(n)$  for some constant  $c < 1$ .
- Solution:**  $T(n) \in \Theta(f(n))$ .

Fall 2009

CS 2233 Discr. Math. Structures

4

## Examples

**Ex.**  $T(n) = 4T(n/2) + n$

$a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n$ .  
**CASE 1:**  $f(n) \in O(n^{2-\varepsilon})$  for  $\varepsilon = 1$ .  
 $\therefore T(n) \in \Theta(n^2)$ .

**Ex.**  $T(n) = 4T(n/2) + n^2$

$a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n^2$ .  
**CASE 2:**  $f(n) \in \Theta(n^2)$ .  
 $\therefore T(n) \in \Theta(n^2 \log n)$ .

## Examples

**Ex.**  $T(n) = 4T(n/2) + n^3$

$a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n^3$ .

**CASE 3:**  $f(n) \in \Omega(n^{2+\varepsilon})$  for  $\varepsilon = 1$   
**and**  $4(n/2)^3 \leq cn^3$  (reg. cond.) for  $c = 1/2$ .  
 $\therefore T(n) \in \Theta(n^3)$ .

**Ex.**  $T(n) = 4T(n/2) + n^2/\log n$

$a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n^2/\log n$ .  
Master method does not apply.